

The zeta function of \mathfrak{g}_{1357A} counting ideals

1 Presentation

\mathfrak{g}_{1357A} has presentation

$$\left\langle x_1, x_2, x_3, x_4, x_5, x_6, x_7 \mid \begin{array}{l} [x_1, x_2] = x_4, [x_1, x_4] = x_5, [x_1, x_5] = x_7, \\ [x_2, x_3] = x_5, [x_2, x_6] = x_7, [x_3, x_4] = -x_7 \end{array} \right\rangle.$$

\mathfrak{g}_{1357A} has nilpotency class 4.

2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{\mathfrak{g}_{1357A,p}}^{\triangleleft}(s) = \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(3s-4)\zeta_p(5s-5)\zeta_p(7s-6).$$

$\zeta_{\mathfrak{g}_{1357A}}^{\triangleleft}(s)$ is uniform.

3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{\mathfrak{g}_{1357A,p}}^{\triangleleft}(s) \Big|_{p \rightarrow p^{-1}} = -p^{21-19s} \zeta_{\mathfrak{g}_{1357A,p}}^{\triangleleft}(s).$$

4 Abscissa of convergence and order of pole

The abscissa of convergence of $\zeta_{\mathfrak{g}_{1357A}}^{\triangleleft}(s)$ is 4, with a simple pole at $s = 4$.

5 Ghost zeta function

This zeta function is its own ghost.

6 Natural boundary

$\zeta_{\mathfrak{g}_{1357A}}^{\triangleleft}(s)$ has meromorphic continuation to the whole of \mathbb{C} .