

# The zeta function of $H \times_{\mathbb{Z}} H$ counting all subrings

## 1 Presentation

$H \times_{\mathbb{Z}} H$  has presentation

$$\langle x_1, x_2, x_3, x_4, y \mid [x_1, x_3] = y, [x_2, x_4] = y \rangle.$$

$H \times_{\mathbb{Z}} H$  has nilpotency class 2.

## 2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{H \times_{\mathbb{Z}} H, p}(s) = \zeta_p(s) \zeta_p(s-1) \zeta_p(s-3) \zeta_p(3s-4) \zeta_p(3s-6) \zeta_p(3s-7) W(p, p^{-s})$$

where  $W(X, Y)$  is

$$1 + X^2Y + X^4Y^2 + X^5Y^3 + X^6Y^3 - X^5Y^4 - X^6Y^4 - X^7Y^5 - X^9Y^6 \\ - X^{11}Y^7.$$

$\zeta_{H \times_{\mathbb{Z}} H}(s)$  is uniform.

## 3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{H \times_{\mathbb{Z}} H, p}(s) \Big|_{p \rightarrow p^{-1}} = -p^{10-5s} \zeta_{H \times_{\mathbb{Z}} H, p}(s).$$

## 4 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{H \times_{\mathbb{Z}} H}(s)$  is 4, with a simple pole at  $s = 4$ .

## 5 Ghost zeta function

The ghost zeta function is the product over all primes of

$$\zeta_p(s) \zeta_p(s-1) \zeta_p(s-3) \zeta_p(3s-4) \zeta_p(3s-6) \zeta_p(3s-7) W_1(p, p^{-s}) W_2(p, p^{-s})$$

where

$$W_1(X, Y) = 1 + X^2Y + X^4Y^2 + X^6Y^3, \\ W_2(X, Y) = 1 - X^5Y^4.$$

The ghost is friendly.

## 6 Natural boundary

$\zeta_{H \times_{\mathbb{Z}} H}(s)$  has a natural boundary at  $\Re(s) = 2$ , and is of type II.