

The zeta function of $M_3 \times \mathbb{Z}^2$ counting all subrings

1 Presentation

$M_3 \times \mathbb{Z}^2$ has presentation

$$\langle z, x_1, x_2, a_1, a_2, x_3 \mid [z, x_1] = x_2, [z, x_2] = x_3 \rangle.$$

$M_3 \times \mathbb{Z}^2$ has nilpotency class 3.

2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\begin{aligned} \zeta_{M_3 \times \mathbb{Z}^2, p}(s) &= \zeta_p(s) \zeta_p(s-1) \zeta_p(s-2) \zeta_p(s-3) \zeta_p(2s-5) \zeta_p(3s-7) \zeta_p(3s-8) \\ &\quad \times \zeta_p(4s-10) W(p, p^{-s}) \end{aligned}$$

where $W(X, Y)$ is

$$\begin{aligned} &1 + X^4 Y^2 + X^5 Y^2 - X^5 Y^3 - X^7 Y^4 + X^{10} Y^4 - 2X^{10} Y^5 - 2X^{11} Y^5 + X^{11} Y^6 \\ &- X^{14} Y^6 - X^{16} Y^7 + X^{16} Y^8 + X^{17} Y^8 + X^{21} Y^{10}. \end{aligned}$$

$\zeta_{M_3 \times \mathbb{Z}^2}(s)$ is uniform.

3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{M_3 \times \mathbb{Z}^2, p}(s) \Big|_{p \rightarrow p^{-1}} = p^{15-6s} \zeta_{M_3 \times \mathbb{Z}^2, p}(s).$$

4 Abscissa of convergence and order of pole

The abscissa of convergence of $\zeta_{M_3 \times \mathbb{Z}^2}(s)$ is 4, with a simple pole at $s = 4$.

5 Ghost zeta function

The ghost zeta function is the product over all primes of

$$\begin{aligned} &\zeta_p(s) \zeta_p(s-1) \zeta_p(s-2) \zeta_p(s-3) \zeta_p(2s-5) \zeta_p(3s-7) \zeta_p(3s-8) \zeta_p(4s-10) \\ &\quad \times W_1(p, p^{-s}) W_2(p, p^{-s}) W_3(p, p^{-s}) \end{aligned}$$

where

$$W_1(X, Y) = 1 + X^5 Y^2 + X^{10} Y^4,$$

$$W_2(X, Y) = 1 - X^4 Y^2 - X^6 Y^3,$$

$$W_3(X, Y) = -1 + X^5 Y^3.$$

The ghost is unfriendly.

6 Natural boundary

$\zeta_{M_3 \times \mathbb{Z}^2}(s)$ has a natural boundary at $\Re(s) = 5/2$, and is of type II.