

The zeta function of $M_3 \times_{\mathbb{Z}} M_3$ counting ideals

1 Presentation

$M_3 \times_{\mathbb{Z}} M_3$ has presentation

$$\langle z_1, z_2, w_1, w_2, x_1, x_2, y \mid [z_1, w_1] = x_1, [z_2, w_2] = x_2, [z_1, x_1] = y, [z_2, x_2] = y \rangle.$$

$M_3 \times_{\mathbb{Z}} M_3$ has nilpotency class 3.

2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{M_3 \times_{\mathbb{Z}} M_3, p}^{\triangleleft}(s) = \zeta_p(s) \zeta_p(s-1) \zeta_p(s-2) \zeta_p(s-3) \zeta_p(3s-4)^2 \zeta_p(5s-5) \zeta_p(7s-4) \\ \times \zeta_p(8s-5) \zeta_p(9s-6) \zeta_p(12s-10) W(p, p^{-s})$$

where $W(X, Y)$ is

$$1 - X^4 Y^5 - 2X^4 Y^8 + X^5 Y^8 + X^4 Y^9 - 2X^5 Y^9 + X^8 Y^{12} - 2X^9 Y^{12} + 3X^9 Y^{13} \\ - 2X^{10} Y^{13} + X^{10} Y^{14} + X^9 Y^{17} + X^{14} Y^{17} + X^{13} Y^{20} - 2X^{13} Y^{21} + 3X^{14} Y^{21} \\ - 2X^{14} Y^{22} + X^{15} Y^{22} - 2X^{18} Y^{25} + X^{19} Y^{25} + X^{18} Y^{26} - 2X^{19} Y^{26} \\ - X^{19} Y^{29} + X^{23} Y^{34}.$$

$\zeta_{M_3 \times_{\mathbb{Z}} M_3}^{\triangleleft}(s)$ is uniform.

3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{M_3 \times_{\mathbb{Z}} M_3, p}^{\triangleleft}(s) \Big|_{p \rightarrow p^{-1}} = -p^{21-17s} \zeta_{M_3 \times_{\mathbb{Z}} M_3, p}^{\triangleleft}(s).$$

4 Abscissa of convergence and order of pole

The abscissa of convergence of $\zeta_{M_3 \times_{\mathbb{Z}} M_3}^{\triangleleft}(s)$ is 4, with a simple pole at $s = 4$.

5 Ghost zeta function

The ghost zeta function is the product over all primes of

$$\zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(3s-4)^2\zeta_p(5s-5)\zeta_p(7s-4)\zeta_p(8s-5) \\ \times \zeta_p(9s-6)\zeta_p(12s-10)W_1(p,p^{-s})W_2(p,p^{-s})W_3(p,p^{-s})$$

where

$$W_1(X, Y) = 1 + X^{14}Y^{17},$$

$$W_2(X, Y) = 1 + X^5Y^8,$$

$$W_3(X, Y) = 1 + X^4Y^9.$$

The ghost is friendly.

6 Natural boundary

$\zeta_{M_3 \times_{\mathbb{Z}} M_3}^{\triangleleft}(s)$ has a natural boundary at $\Re(s) = 14/17$, and is of type III.

7 Notes

This ideal zeta function is identical to that of \mathfrak{g}_{137B} , though the Lie rings themselves are non-isomorphic.