

# The zeta function of $T_4$ counting all subrings

## 1 Presentation

$T_4$  has presentation

$$\langle x_1, x_2, x_3, x_4, y_1, y_2, y_3 \mid [x_1, x_2] = y_1, [x_2, x_3] = y_2, [x_3, x_4] = y_3 \rangle.$$

$T_4$  has nilpotency class 2.

## 2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{T_4, p}(s) = \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(2s-5)^2\zeta_p(2s-6)^2\zeta_p(3s-6) \\ \times \zeta_p(3s-8)^2\zeta_p(3s-9)\zeta_p(4s-12)\zeta_p(5s-14)W(p, p^{-s})$$

where  $W(X, Y)$  is

$$\begin{aligned} & 1 + X^4Y^2 + X^5Y^2 - 2X^5Y^3 - 3X^6Y^3 + X^6Y^4 - X^8Y^4 - 2X^9Y^4 - 2X^{10}Y^4 \\ & + X^{11}Y^5 - 2X^{12}Y^5 - 4X^{13}Y^5 - X^{14}Y^5 + X^{11}Y^6 + X^{12}Y^6 + 4X^{13}Y^6 \\ & + 2X^{14}Y^6 + 2X^{15}Y^6 - 2X^{16}Y^6 - 2X^{17}Y^6 + X^{14}Y^7 + 3X^{15}Y^7 + 2X^{16}Y^7 \\ & + 4X^{17}Y^7 + 2X^{18}Y^7 - X^{19}Y^7 + X^{20}Y^7 - X^{15}Y^8 + 3X^{18}Y^8 + 6X^{19}Y^8 \\ & + 4X^{20}Y^8 + 3X^{21}Y^8 - X^{18}Y^9 - 4X^{19}Y^9 - 4X^{20}Y^9 - 2X^{21}Y^9 + 5X^{22}Y^9 \\ & + 2X^{23}Y^9 + 4X^{24}Y^9 + 2X^{25}Y^9 - X^{20}Y^{10} - 3X^{22}Y^{10} - 8X^{23}Y^{10} \\ & - 3X^{24}Y^{10} - X^{25}Y^{10} + 2X^{26}Y^{10} + 2X^{27}Y^{10} + 2X^{23}Y^{11} - 2X^{24}Y^{11} \\ & - 4X^{25}Y^{11} - 8X^{26}Y^{11} - 11X^{27}Y^{11} - 4X^{28}Y^{11} + X^{29}Y^{11} + 3X^{30}Y^{11} \\ & + X^{25}Y^{12} + 4X^{26}Y^{12} + 3X^{27}Y^{12} + 2X^{28}Y^{12} - 6X^{29}Y^{12} - 11X^{30}Y^{12} \\ & - 6X^{31}Y^{12} - 4X^{32}Y^{12} + X^{33}Y^{12} + 2X^{29}Y^{13} + 6X^{30}Y^{13} + 5X^{31}Y^{13} \\ & - 5X^{33}Y^{13} - 6X^{34}Y^{13} - 2X^{35}Y^{13} - X^{31}Y^{14} + 4X^{32}Y^{14} + 6X^{33}Y^{14} \\ & + 11X^{34}Y^{14} + 6X^{35}Y^{14} - 2X^{36}Y^{14} - 3X^{37}Y^{14} - 4X^{38}Y^{14} - X^{39}Y^{14} \\ & - 3X^{34}Y^{15} - X^{35}Y^{15} + 4X^{36}Y^{15} + 11X^{37}Y^{15} + 8X^{38}Y^{15} + 4X^{39}Y^{15} \\ & + 2X^{40}Y^{15} - 2X^{41}Y^{15} - 2X^{37}Y^{16} - 2X^{38}Y^{16} + X^{39}Y^{16} + 3X^{40}Y^{16} \\ & + 8X^{41}Y^{16} + 3X^{42}Y^{16} + X^{44}Y^{16} - 2X^{39}Y^{17} - 4X^{40}Y^{17} - 2X^{41}Y^{17} \\ & - 5X^{42}Y^{17} + 2X^{43}Y^{17} + 4X^{44}Y^{17} + 4X^{45}Y^{17} + X^{46}Y^{17} - 3X^{43}Y^{18} \\ & - 4X^{44}Y^{18} - 6X^{45}Y^{18} - 3X^{46}Y^{18} + X^{49}Y^{18} - X^{44}Y^{19} + X^{45}Y^{19} \\ & - 2X^{46}Y^{19} - 4X^{47}Y^{19} - 2X^{48}Y^{19} - 3X^{49}Y^{19} - X^{50}Y^{19} + 2X^{47}Y^{20} \end{aligned}$$

$$\begin{aligned}
& + 2X^{48}Y^{20} - 2X^{49}Y^{20} - 2X^{50}Y^{20} - 4X^{51}Y^{20} - X^{52}Y^{20} - X^{53}Y^{20} \\
& + X^{50}Y^{21} + 4X^{51}Y^{21} + 2X^{52}Y^{21} - X^{53}Y^{21} + 2X^{54}Y^{22} + 2X^{55}Y^{22} \\
& + X^{56}Y^{22} - X^{58}Y^{22} + 3X^{58}Y^{23} + 2X^{59}Y^{23} - X^{59}Y^{24} - X^{60}Y^{24} - X^{64}Y^{26}.
\end{aligned}$$

$\zeta_{T_4}(s)$  is uniform.

### 3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{T_4,p}(s)|_{p \rightarrow p^{-1}} = -p^{21-7s}\zeta_{T_4,p}(s).$$

### 4 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{T_4}(s)$  is 4, with a simple pole at  $s = 4$ .

### 5 Ghost zeta function

The ghost zeta function is the product over all primes of

$$\begin{aligned}
& \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(2s-5)^2\zeta_p(2s-6)^2\zeta_p(3s-6)\zeta_p(3s-8)^2 \\
& \times \zeta_p(3s-9)\zeta_p(4s-12)\zeta_p(5s-14)W_1(p,p^{-s})W_2(p,p^{-s})W_3(p,p^{-s}) \\
& \times W_4(p,p^{-s})W_5(p,p^{-s})
\end{aligned}$$

where

$$\begin{aligned}
W_1(X, Y) &= 1 + X^{20}Y^7, \\
W_2(X, Y) &= 1 - X^{19}Y^7, \\
W_3(X, Y) &= -1 + X^5Y^2 + X^{10}Y^4, \\
W_4(X, Y) &= 1 - X^9Y^4, \\
W_5(X, Y) &= -1 - X^6Y^4.
\end{aligned}$$

The ghost is unfriendly.

### 6 Natural boundary

$\zeta_{T_4}(s)$  has a natural boundary at  $\Re(s) = 20/7$ , and is of type III.