

The zeta function of $\mathfrak{g}_{6,16}$ counting ideals

1 Presentation

$\mathfrak{g}_{6,16}$ has presentation

$$\left\langle x_1, x_2, x_3, x_4, x_5, x_6 \mid \begin{array}{l} [x_1, x_3] = x_4, [x_1, x_4] = x_5, [x_1, x_5] = x_6, \\ [x_2, x_3] = x_5, [x_2, x_4] = x_6 \end{array} \right\rangle.$$

$\mathfrak{g}_{6,16}$ has nilpotency class 4.

2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{\mathfrak{g}_{6,16},p}^{\triangleleft}(s) = \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(3s-3)\zeta_p(5s-4)\zeta_p(6s-3)\zeta_p(7s-5) \\ \times \zeta_p(7s-3)^{-1}.$$

$\zeta_{\mathfrak{g}_{6,16}}^{\triangleleft}(s)$ is uniform.

3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{\mathfrak{g}_{6,16},p}^{\triangleleft}(s) \Big|_{p \rightarrow p^{-1}} = p^{15-17s} \zeta_{\mathfrak{g}_{6,16},p}^{\triangleleft}(s).$$

4 Abscissa of convergence and order of pole

The abscissa of convergence of $\zeta_{\mathfrak{g}_{6,16}}^{\triangleleft}(s)$ is 3, with a simple pole at $s = 3$.

5 Ghost zeta function

This zeta function is its own ghost.

6 Natural boundary

$\zeta_{\mathfrak{g}_{6,16}}^{\triangleleft}(s)$ has meromorphic continuation to the whole of \mathbb{C} .