

The zeta function of $\mathfrak{g}_{6,4}$ counting all subrings

1 Presentation

$\mathfrak{g}_{6,4}$ has presentation

$$\langle x_1, x_2, x_3, x_4, x_5, x_6 \mid [x_1, x_2] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = x_6 \rangle.$$

$\mathfrak{g}_{6,4}$ has nilpotency class 2.

2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\begin{aligned} \zeta_{\mathfrak{g}_{6,4},p}(s) &= \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(2s-5)\zeta_p(3s-5)\zeta_p(3s-7) \\ &\quad \times \zeta_p(3s-8)\zeta_p(4s-9)\zeta_p(4s-11)\zeta_p(5s-12)W(p, p^{-s}) \end{aligned}$$

where $W(X, Y)$ is

$$\begin{aligned} &1 + X^4Y^2 - X^5Y^3 + X^6Y^3 - X^6Y^4 - X^7Y^4 - X^9Y^4 + X^{10}Y^4 - X^9Y^5 \\ &- 2X^{10}Y^5 - 3X^{11}Y^5 - 2X^{12}Y^5 - X^{13}Y^5 + X^{10}Y^6 + X^{11}Y^6 + 2X^{12}Y^6 \\ &+ X^{13}Y^6 + X^{14}Y^6 - X^{15}Y^6 - X^{13}Y^7 - X^{14}Y^7 - 2X^{15}Y^7 - X^{16}Y^7 \\ &- X^{17}Y^7 + X^{14}Y^8 + 2X^{15}Y^8 + 3X^{16}Y^8 + 3X^{17}Y^8 + X^{18}Y^8 - X^{19}Y^8 \\ &+ X^{20}Y^8 - X^{17}Y^9 + X^{18}Y^9 + 2X^{19}Y^9 + 2X^{20}Y^9 + 2X^{21}Y^9 + X^{22}Y^9 \\ &+ X^{22}Y^{10} + X^{23}Y^{10} + X^{24}Y^{10} - X^{21}Y^{11} - X^{22}Y^{11} + X^{26}Y^{11} + X^{27}Y^{11} \\ &- X^{24}Y^{12} - X^{25}Y^{12} - X^{26}Y^{12} - X^{26}Y^{13} - 2X^{27}Y^{13} - 2X^{28}Y^{13} \\ &- 2X^{29}Y^{13} - X^{30}Y^{13} + X^{31}Y^{13} - X^{28}Y^{14} + X^{29}Y^{14} - X^{30}Y^{14} - 3X^{31}Y^{14} \\ &- 3X^{32}Y^{14} - 2X^{33}Y^{14} - X^{34}Y^{14} + X^{31}Y^{15} + X^{32}Y^{15} + 2X^{33}Y^{15} \\ &+ X^{34}Y^{15} + X^{35}Y^{15} + X^{33}Y^{16} - X^{34}Y^{16} - X^{35}Y^{16} - 2X^{36}Y^{16} - X^{37}Y^{16} \\ &- X^{38}Y^{16} + X^{35}Y^{17} + 2X^{36}Y^{17} + 3X^{37}Y^{17} + 2X^{38}Y^{17} + X^{39}Y^{17} \\ &- X^{38}Y^{18} + X^{39}Y^{18} + X^{41}Y^{18} + X^{42}Y^{18} - X^{42}Y^{19} + X^{43}Y^{19} - X^{44}Y^{20} \\ &- X^{48}Y^{22}. \end{aligned}$$

$\zeta_{\mathfrak{g}_{6,4}}(s)$ is uniform.

3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{\mathfrak{g}_{6,4},p}(s) \Big|_{p \rightarrow p^{-1}} = p^{15-6s} \zeta_{\mathfrak{g}_{6,4},p}(s).$$

4 Abscissa of convergence and order of pole

The abscissa of convergence of $\zeta_{\mathfrak{g}_{6,4}}(s)$ is 4, with a simple pole at $s = 4$.

5 Ghost zeta function

The ghost zeta function is the product over all primes of

$$\begin{aligned} & \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(2s-5)\zeta_p(3s-5)\zeta_p(3s-7)\zeta_p(3s-8) \\ & \times \zeta_p(4s-9)\zeta_p(4s-11)\zeta_p(5s-12)W_1(p,p^{-s})W_2(p,p^{-s})W_3(p,p^{-s}) \\ & \times W_4(p,p^{-s}) \end{aligned}$$

where

$$\begin{aligned} W_1(X, Y) &= 1 - X^{13}Y^5, \\ W_2(X, Y) &= -1 + X^7Y^3 + X^{14}Y^6 - X^{21}Y^9, \\ W_3(X, Y) &= -1 - X^4Y^2 + X^8Y^4, \\ W_4(X, Y) &= 1 - X^6Y^4. \end{aligned}$$

The ghost is unfriendly.

6 Natural boundary

$\zeta_{\mathfrak{g}_{6,4}}(s)$ has a natural boundary at $\Re(s) = 13/5$, and is of type III.

7 Notes

This zeta function is a lot more complicated than the one only counting ideals!

The Lie ring is sometimes written as $(F_{2,3}/\langle z \rangle) \cdot \mathbb{Z}$.