

# The zeta function of $\mathfrak{p}_3$ counting all subgroups

## 1 Presentation

$\mathfrak{p}_3$  has presentation

$$\langle x, y, r \mid [x, y], r^3, x^r = x^{-1}y, y^r = x^{-1} \rangle.$$

## 2 The zeta function itself

The zeta function was calculated by du Sautoy, McDermott and Smith. It is

$$\zeta_{\mathfrak{p}_3}(s) = \zeta(s-1)L(s-1, \chi_3) + 3^{-s}\zeta(s)\zeta(s-1),$$

where  $\chi_3$  is defined by

$$\chi_3(a) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{3} \\ -1 & \text{if } a \equiv -1 \pmod{3} \\ 0 & \text{otherwise} \end{cases},$$

and  $L(s, \chi_3)$  is the Dirichlet  $L$ -function of  $\chi_3$ ,

$$L(s, \chi_3) = \sum_{n=1}^{\infty} \chi_3(n)n^{-s}.$$

## 3 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{\mathfrak{p}_3}(s)$  is 2, with a simple pole at  $s = 2$ . Since this group is a finite extension of a free abelian group, its zeta function has meromorphic continuation to  $\mathbb{C}$ .