

# The zeta function of $\mathbf{p3m1}$ counting all subgroups

## 1 Presentation

$\mathbf{p3m1}$  has presentation

$$\left\langle x, y, r, m \mid \begin{array}{l} [x, y], r^3, m^2, r^m = r^{-1}, x^r = x^{-1}y, \\ y^r = x^{-1}, x^m = x^{-1}, y^m = x^{-1}y \end{array} \right\rangle.$$

## 2 The zeta function itself

The zeta function was calculated by du Sautoy, McDermott and Smith. It is

$$\begin{aligned} \zeta_{\mathbf{p3m1}}(s) &= (1 + 9 \cdot 3^{-s})\zeta(2s - 2) + 2^{-s}\zeta(s - 1)L(s - 1, \chi_3) \\ &\quad + (3 \cdot 3^{-s} - 2 \cdot 6^{-s} + 12 \cdot 12^{-s})\zeta(s)\zeta(s - 1), \end{aligned}$$

where  $\chi_3$  is defined by

$$\chi_3(a) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{3} \\ -1 & \text{if } a \equiv -1 \pmod{3}, \\ 0 & \text{otherwise} \end{cases}$$

and  $L(s, \chi_3)$  is the Dirichlet  $L$ -function of  $\chi_3$ ,

$$L(s, \chi_3) = \sum_{n=1}^{\infty} \chi_3(n)n^{-s}.$$

## 3 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{\mathbf{p3m1}}(s)$  is 2, with a simple pole at  $s = 2$ . Since this group is a finite extension of a free abelian group, its zeta function has meromorphic continuation to  $\mathbb{C}$ .